

**Bonnie P. Gregory, MD**  
**GLENOID RECONSTRUCTION**  
**REHABILITATION PROTOCOL**

	<b>RANGE OF MOTION</b>	<b>IMMOBILIZER</b>	<b>EXERCISES</b>
<i>PHASE I</i> <i>0-6 weeks</i>	Limit ER to passive 45 ° to protect subscap repair FE progress as tolerated	0-2 weeks: Worn at all times (day and night) Off for gentle exercise only 2-6 weeks: Worn daytime only	0-3 weeks: Grip strengthening, pendulum exercises Elbow/wrist/hand ROM at home 3-6 weeks: Begin cuff, deltoid isometrics; limit ER to passive 45 ° No active IR nor extension until 6 weeks
<i>PHASE II</i> <i>6-12 weeks</i>	Increase as tolerated to full Begin active assisted/active internal rotation and extension as tolerated after 6 weeks	None	6-8 weeks: Begin light resisted ER, forward flexion and abduction 8-12 weeks: Begin resisted internal rotation, extension and scapular retraction
<i>PHASE III</i> <i>12-24 weeks</i>	Progress to full motion without discomfort	None	Advance strengthening as tolerated Closed chain scapular rehab and functional rotator cuff strengthening; focus on anterior deltoid and teres Maximize subscapular stabilization

For any questions or concerns regarding the protocol or rehabilitation process please contact:

Dr. Gregory's Office:  
Phone: 713-486-7080  
Fax: 713-452-4143

<https://www.bonniegregorymd.com/>